

The inner world of overactive monitoring: neural markers of interoception in obsessive–compulsive disorder

A. Yoris^{1,2,3}, A. M. García^{1,2,4}, L. Traiber³, H. Santamaría-García^{1,2,5,6}, M. Martorell¹, F. Alifano¹, R. Kichic^{1,3}, J. S. Moser⁷, M. Cetkovich¹, F. Manes^{1,2,8}, A. Ibáñez^{1,2,8,9,10} and L. Sedeño^{1,2*}

¹Laboratory of Experimental Psychology and Neuroscience (LPEN), Institute of Cognitive and Translational Neuroscience (INCyT), INECO Foundation, Favaloro University, Buenos Aires, Argentina

²National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina

³Anxiety and Trauma Clinic, INECO (Institute of Cognitive Neurology), Buenos Aires, Argentina

⁴Faculty of Education, National University of Cuyo, Mendoza, Argentina

⁵Centro de Memoria y Cognición Intellectus, Hospital Universitario San Ignacio, Bogotá, Colombia

⁶Pontificia Universidad Javeriana, Bogotá, Colombia

⁷Department of Psychology, Michigan State University, East Lansing, MI, USA

⁸Centre of Excellence in Cognition and its Disorders, Australian Research Council (ACR), Sydney, Australia

⁹Universidad Autónoma del Caribe, Barranquilla, Colombia

¹⁰Center for Social and Cognitive Neuroscience (CSCN), School of Psychology, Universidad Adolfo Ibáñez, Santiago de Chile, Chile

Background. Obsessive–compulsive disorder (OCD) patients typically overmonitor their own behavior, as shown by symptoms of excessive doubt and checking. Although this is well established for the patients' relationship with external stimuli in the environment, no study has explored their monitoring of internal body signals, a process known to be affected in anxiety-related syndromes. Here, we explored this issue through a cardiac interoception task that measures sensing of heartbeats. Our aim was to explore key behavioral and electrophysiological aspects of internal-cue monitoring in OCD, while examining their potential distinctiveness in this condition.

Method. We administered a heartbeat detection (HBD) task (with related interoceptive confidence and awareness measures) to three matched groups (OCD patients, panic disorder patients, healthy controls) and recorded ongoing modulations of two task-relevant electrophysiological markers: the heart evoked potential (HEP) and the motor potential (MP).

Results. Behaviorally, OCD patients outperformed controls and panic patients in the HBD task. Moreover, they exhibited greater amplitude modulation of both the HEP and the MP during cardiac interoception. However, they evinced poorer confidence and awareness of their interoceptive skills.

Conclusions. Convergent behavioral and electrophysiological data showed that overactive monitoring in OCD extends to the sensing of internal bodily signals. Moreover, this pattern discriminated OCD from panic patients, suggesting a condition-distinctive alteration. Our results highlight the potential of exploring interoceptive processes in the OCD spectrum to better characterize the population's cognitive profile. Finally, these findings may lay new bridges between somatic theories of emotion and cognitive models of OCD.

Received 14 April 2016; Revised 25 January 2017; Accepted 26 January 2017; First published online 4 April 2017

Key words: Heart evoked potential, interoception, motor potential, obsessive–compulsive disorder, overactive monitoring.

Introduction

Obsessive–compulsive disorder (OCD) is characterized by intrusive distressing ideas, repeated behaviors and neutralizing thoughts, which aim to control internal and external events (Aouizerate *et al.* 2004). In this

context, OCD patients typically develop overactive self-performance monitoring (Endrass & Ullsperger, 2014). Such an alteration is accompanied by event-related potential (ERP) abnormalities in components indexing monitoring processes, such as the error-related negativity (Gehring *et al.* 2000; Riesel *et al.* 2014), the correct-response negativity (CRN) (Riesel *et al.* 2011; Carrasco *et al.* 2013), and the motor potential (MP) (Johannes *et al.* 2001). Note that the latter potential, which is elicited by voluntary finger or foot movements (Brunia & Van den Bosch, 1984; Deecke *et al.* 1984), exhibits larger amplitudes for correct than

* Address for correspondence: L. Sedeño, Laboratory of Experimental Psychology and Neuroscience (LPEN), Institute of Cognitive and Translational Neuroscience (INCyT), C1078AAI, Pacheco de Melo 1860, Buenos Aires, Argentina.
(Email: lucas.sedeno@gmail.com)

incorrect responses, thus yielding information about monitoring processes (Vaughan *et al.* 1968; Deecke *et al.* 1984; Simons, 2010). While overactive monitoring is well established in OCD (Gehring *et al.* 2000; Endrass & Ullsperger, 2014) and may even represent an endophenotype of the condition (Riesel *et al.* 2011, 2015), little is known about the specificity and scope of this pattern. Only a few studies have explored it by comparing OCD and other psychiatric conditions (e.g. Xiao *et al.* 2011), and none has examined it in relation to inwards attention. Here we address both issues combining behavioral and neurophysiological measures.

Available findings stem from responses to external stimuli. However, given the role of peripheral physiological processes in homeostasis and emotional regulation (Cameron, 2001; Craig, 2002; Weiss *et al.* 2014), OCD may also involve overactive monitoring of internal signals, such as those indexing bodily states. A crucial function in this respect is interoception, the perceptual and physiological sensing of visceral signals (Craig, 2003). This multidimensional function (Craig, 2002; Garfinkel *et al.* 2015; García-Cordero *et al.* 2016) differentially engages the insula, the anterior cingulate cortex and the prefrontal cortex (Critchley *et al.* 2004; Zaki *et al.* 2012). Notably, the former two areas evidence increased activity and volume in OCD patients (Song *et al.* 2011). Thus, interoceptive alterations may be reasonably expected in this population.

Critical evidence on bodily signal monitoring is afforded by heartbeat detection (HBD) tasks, in which participants follow their own heartbeats through mental (Schandry *et al.* 1993) or motor (Couto *et al.* 2014) tracking. These tasks tap interoceptive accuracy (IAcc) – the participants' objective precision in detecting visceral sensations (Garfinkel *et al.* 2015) –, and they also allow assessment of two related dimensions: interoceptive sensibility (ISen) – the subject's confidence in his/her own performance – and interoceptive awareness (IAw) – a metacognitive index of interoceptive performance (Zaki *et al.* 2012; Garfinkel *et al.* 2015). Together with the evidence above, reports of metacognitive impairments in OCD (Cucchi *et al.* 2012; Coles *et al.* 2015) suggest that such three interoceptive dimensions may be altered in this group.

Finally, electrophysiological correlates of heartbeat sensing are indexed by the heart evoked potential (HEP). This ERP deflection reflects attention allocation to cardiac modulations irrespective of response accuracy (Schandry *et al.* 1986; Pollatos & Schandry, 2004). If overactive monitoring in OCD extends to the interoceptive domain, then distinct HEP modulations should also be observed during the HBD task.

Our study examined whether overactive monitoring in OCD is also present in the interoceptive domain,

and whether this potential pattern is different in other clinical conditions, with a specific focus on panic disorder. To this end, we compared the performance of three samples (OCD patients, panic disorder patients and healthy controls) on a motor-tracking HBD task. We measured IAcc, ISen and IAw, together with associated neurophysiological processes. Since the HBD task does not include high-conflict trials but requires precise manual responses to cardiac signals, our analysis focused on the MP and the HEP rather than the ERN. These two components offer complementary measures of attention allocation to inner-body modulations. Specifically, we hypothesized that, relative to both other samples, OCD patients would exhibit: (i) enhanced IAcc and impaired IAw; (ii) increased MP amplitude for correct relative to incorrect trials; and (iii) enlarged HEP modulations, indexing overactive monitoring of internal signals. Our study thus offers an unprecedented cross-dimensional view into inner monitoring in OCD. Insights derived therefrom could contribute to: (a) a better characterization of the patients' cognitive profile and the relation between inner monitoring and obsessive-compulsive mechanisms and symptoms; (b) the formalization of potentially sensitive and specific biomarkers of the disease; and (c) the establishment of new avenues to assess disease progression and response to clinical treatment.

Method

Participants

The study comprised 15 OCD patients, 15 panic disorder patients and 25 healthy volunteers. The patients' diagnoses were established by two OCD experts (R.K. and A.Y.) using the Structured Clinical Interview for DSM-IV Axis I SCID-I (First *et al.* 2002) and following Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) criteria (American Psychiatric Association, 1994). OCD patients scored well above the 21-point cut-off (mean = 30.27) on the Obsessive-Compulsive Inventory (OCI-R) (Foa *et al.* 2002), and they presented high scores (mean = 80.93) in the Obsessive Belief Questionnaire (OBQ-TRIP) (Moulding *et al.* 2011) (see online Supplementary Table S1.1). OCD and panic patients reported psychiatric medication intake during recruitment (see online Supplementary Table S1.2). Healthy controls had no history of drug abuse or neuropsychiatric disease. The groups were matched for age, gender, education, handedness and body mass index (Table 1). Participants provided informed consent in accordance with the Declaration of Helsinki. The study was approved by the institutional ethics committee.

Table 1. Demographic and mood/anxiety results

Variables	Group			Statistics	Post-hoc analyses (Tukey)		
	OCD	CTRL	PANIC		OCD v. CTRL	CTRL v. PANIC	OCD v. PANIC
Gender, <i>n</i>				$\chi^2 = 6.35, p = 0.09$	–	–	–
Female	8	14	9				
Male	7	11	6				
Age, years	30.40 (9.98)	33.45 (11.55)	35.46 (8.99)	$F = 1.58, p = 0.21,$ $\eta p^2 = 0.06$	–	–	–
Education	15.58 (2.06)	16.19 (2.93)	17.50 (2.89)	$F = 1.12, p = 0.29,$ $\eta p^2 = 0.03$	–	–	–
Body mass index, kg/m ²	23.52 (3.32)	23.35 (3.43)	24.79 (3.84)	$F = 0.65, p = 0.52,$ $\eta p^2 = 0.06$	–	–	–
Handedness, <i>n</i>							
Right	15	25	15				
Left	0	0	0				
HADS-A	13.66 (1.49)	6.20 (2.64)	11.46 (3.64)	$F = 12.70, p < 0.01^*,$ $\eta p^2 = 0.35$	$p < 0.01^*$	$p < 0.01^*$	$p = 0.2$
HADS-D	9.80 (3.84)	2.65 (2.87)	7.46 (4.61)	$F = 16.73, p < 0.01^*,$ $\eta p^2 = 0.41$	$p < 0.01^*$	$p < 0.01^*$	$p < 0.01$
STAI-S	32.73 (7.21)	32.4 (7.05)	35.90 (7.56)	$F = 0.85, p = 0.43,$ $\eta p^2 = 0.03$	–	–	–

Data are given as mean (standard deviation) unless otherwise indicated.

OCD, Obsessive–compulsive disorder; CTRL, control; PANIC, panic; HADS-A, Hospital Anxiety and Depression Scale, anxiety subscale; HADS-D, Hospital Anxiety and Depression Scale, depression subscale; STAI-S, State Trait Anxiety Inventory, state version.

* Significant difference.

Mood and anxiety measures

Anxiety and depression symptoms were evaluated using the Hospital Anxiety and Depression Scale (HADS). The HADS has two subscales, addressing recent depressive symptoms (HADS-D) and trait anxiety (HADS-A) (Zigmond & Snaith, 1983). Anxiety levels during the HBD task were assessed via the state version of the State Trait Anxiety Inventory (STAI-S; Spielberger *et al.* 1970).

HBD task

Cardiac interoception was assessed through a modified version of a validated HBD task (Sedeno *et al.* 2014; Couto *et al.* 2015). Two conditions were considered, both involving dominant-hand responses. In the exteroceptive condition, participants tapped a keyboard to follow an audio-recording of a typical heartbeat; this indexed monitoring of external stimuli. In the interoceptive condition, participants followed their own heartbeats without any external cues, which provided a measure of IAcc (Garfinkel *et al.* 2015) (see online Supplementary material section 2 for further details). In addition, we assessed ISen and

IAw (Garfinkel *et al.* 2015), as defined in the Introduction. Finally, during this task, neurophysiological and cardiac signals were recorded via electroencephalography (EEG) and electrocardiography (ECG).

IAcc

Behavioral interoceptive performance was indexed by tapping accuracy (TA) (Canales-Johnson *et al.* 2015). First, correct answers were selected based on responses that were temporally locked to the R-peak of the heartbeats. We used this R-wave deflection given that the time window between 200 to 300 ms after it is a critical proxy of heartbeat perception (Brenner & Kluitse, 1988; Wiens & Palmer, 2001; Critchley *et al.* 2004; Kleckner *et al.* 2015).

TA was defined as the average of the absolute value of the time difference between the R-peak and the motor tap in correct answers (Canales-Johnson *et al.* 2015), namely: $TA = \sum(\text{motor tap time} - \text{R-peak time})/n$. Accordingly, given that TA is a reaction-time measure, smaller values indicate better behavioral performance (IAcc) during the HBD task.

ISen

ISen represents a subject's self-confidence relative to his/her objective performance in an interoceptive task (Garfinkel et al. 2015). To measure this variable, at the end of the interoceptive condition of the HBD task, we asked participants to rate their confidence in their performance from 1 (not confident at all) to 9 (fully confident).

IAw

IAw represents the degree to which objective performance during the HBD task (IAcc) is associated with subjective confidence in such performance (ISen) (Garfinkel et al. 2015). This index was calculated for each group (given that we only had one measure for IAcc and ISen per participant) based on an indirect correlation between IAcc (based on TA) and ISen. A high correspondence between these two measures, represented by a high correlation value, indicates the ability of participants to recognize that their HBD performance was either good or bad (Garfinkel et al. 2015).

ERP measures*EEG recordings and preprocessing steps*

EEG signals were recorded with a Biosemi Active-two 128-channel system at 1024 Hz, resampled offline at 256 Hz. The reference was set by default to link mastoids, and re-referenced off-line to the average electrodes. Segments with eye movement contamination were removed from further analysis using independent component analysis (ICA) and a visual inspection procedure. To control for potential cardiac artifacts, we also removed cardiac components via ICA. Two external electrodes were used to collect the ECG signal. R-wave-ECG detection was achieved by a peak finder function implemented on MATLAB (MathWorks, Inc., USA), which quickly identifies local peaks or valleys (local extrema) in a noisy vector using the alternating nature of the derivatives and a user-defined magnitude threshold to determine whether each peak is significantly larger (or smaller) than the data around it (Kruczyk et al. 2013). See online Supplementary material section 3 for EEG preprocessing details.

HEP

The HEP is a modulation emerging 200–500 ms after the R-wave peak (Montoya et al. 1993; Pollatos & Schandry, 2004; Fukushima et al. 2011; Canales-Johnson et al. 2015) that indexes attention to cardiac modulations (Schandry & Weitkunat, 1990; Montoya et al. 1993), and also other processes related to body-brain communication, such as emotional processes (Canales-Johnson et al. 2015), motivation (Schandry &

Weitkunat, 1990), stress (Gray et al. 2007) and pain perception (Shao et al. 2011). In all of these, HEP amplitude seems to be triggered by attention to body sensations.

Regarding cardiac perception, one main pathway of heartbeat information relies on signals from baroreceptors in the aortic arch and carotid bodies (Critchley et al. 2004; Critchley & Harrison, 2013) which, primarily via the vagus nerve and its connections to the nucleus of the solitary tracks (Craig, 2002, Janig, 1996), provide projections to centers such as the hypothalamus, thalamus and cerebral cortex (Craig, 2002). Here, the insular cortex and the ACC play a major role as a viscerosensory (Craig, 2002, 2003) and visceromotor centers, respectively (Craig, 2002, 2004; Dum et al. 2009). Primary and secondary somatosensory cortices constitute another pathway of cardiac information, given that sensory information from the skin of the chest is also essential for interoceptive perception (Khalsa et al. 2009; Couto et al. 2014). The key role of these cortical areas (insula, ACC and somatosensory cortices) has been supported by lesion models (Khalsa et al. 2009; Couto et al. 2014; Ronchi et al. 2015; Terasawa et al. 2015; García-Cordero et al. 2016), fMRI studies and meta-analyses (Critchley et al. 2004; Pollatos et al. 2007; Schulz et al. 2016), and brain stimulation studies (Pollatos et al. 2016). Critically, HEP source analysis has also revealed the involvement of these areas in cardiac perception (Pollatos et al. 2005), which supports the relationship between this ERP and interoceptive processing.

Based on previous studies (Pollatos & Schandry, 2004; Gray et al. 2007), we measured this potential in a region of interest (ROI) comprising four right-prefrontal electrodes (C14, C15, C16, C10). A baseline was defined at 200 ms before the R-peak and the epochs were selected between -0.2 and 0.5 s. We predicted greater amplitude of the HEP for internally than externally triggered cardiac stimuli, and for OCD patients in general.

MP

The MP waveform begins 150 to 100 ms prior to the onset of a self-paced movement, although it can peak 50 ms after the movement (Deecke et al. 1980). Following standard recommendations for MP research (Johannes et al. 2001) and previous protocols (Caldara et al. 2004), we selected six motor-related electrodes to construct a left-hemisphere ROI, considering that all subjects were right-handed (D11, D12, D13, D18, D19, D20). The MP can be a promising ERP for interoceptive performance, given that interoceptive processes are linked to insular-motor input of internal and external stimuli (Nagai et al. 2007). Recent findings have shown cortical alterations in the insula and related

structures in OCD (Song *et al.* 2011). Furthermore, ERP analysis comparing OCD and panic patients have shown similar inhibitory deficits in early potentials associated to motor response during a Go/NoGo task (Bannon *et al.* 2002). Building on previous research (Caldara *et al.* 2004), we analysed intra-group variance comparing right and wrong answers. We assumed that correct responses should be affected by higher effort or monitoring, while incorrect responses should not show such modulations. Specifically, we expected MP amplitude to increase for correct relative to incorrect responses, thus framing it as an indicator of precision and quickness of monitoring functions.

Data analysis

Demographic and clinical data were assessed through analyses of variance. Gender was analysed with the Pearson χ^2 test. Considering the possible influence of depression and anxiety symptoms on interoception (Dunn *et al.* 2007; Domschke *et al.* 2010) and the significant mood differences between groups (Table 1), we performed an analysis of covariance using HADS scores as covariates for the HBD measures (only significant results surviving covariance are reported). Effect sizes were calculated with partial η^2 . *Post-hoc* analyses were performed with Tukey's honestly significant difference test. We also used multiple regression models ($p < 0.05$) to assess the relationship between attention to heartbeats (IAcc) and body signals (HEP) from the HBD task, on the one hand, and the dysfunctional beliefs of OCD patients (perfectionism, inflated responsibility, intolerance of uncertainty, need to control, and overestimation of threat), as measured by the OBQ-TRIP, on the other (see online Supplementary material section 4).

Point-by-point comparisons along the ERP were made via the Monte Carlo permutation test (Manly, 2007), as done in previous HEP (Couto *et al.* 2015) and MP (Ibanez *et al.* 2013) studies. If the ensuing p values are smaller than the critical α -level of 0.05, then the data can be concluded to reveal significant differences. This method circumvents the multiple comparisons problem and does not depend on multiple comparison corrections or Gaussian distribution assumptions (Nichols & Holmes, 2002). In addition, non-*a priori* windows were selected to analyse the ERPs. Importantly, by adopting a data-driven approach in our analyses, we avoided the bias of *a priori* window selection.

Results

Demographic results

No group differences were found in gender, age, formal education, handedness or body mass index (Table 1).

Mood and anxiety results

Data gleaned through the HADS-A revealed more anxiety symptoms in OCD and panic patients than in controls. No differences were found between the patient groups. Results from the HADS-D subscale showed that depression scores were highest in OCD patients, followed by panic patients, and then by controls. No between-group differences were found in the STAI-S (Table 1).

HBD task

IAcc

All groups performed similarly in the exteroceptive condition. However, significant differences emerged in the IAcc measure (Fig. 1a and online Supplementary Table S5.1). Compared with controls and panic patients, OCD patients presented significantly lower values of TA, indicating higher IAcc (i.e. better synchronization of motor responses to heartbeats).

ISen

Analysis of ISen scores revealed significantly lower self-confidence in OCD patients relative to controls ($p = 0.03$) and panic patients ($p = 0.04$), who did not differ from each other ($p = 0.1$) (see online Supplementary Table S5.1).

IAw

Spearman correlations revealed a significant association between IAcc and ISen in controls ($R = -0.46$, $p = 0.02$) and panic patients ($R = -0.55$, $p = 0.03$). Given that IAcc is a reaction time score (TA), the negative direction of the relationship indicates a correspondence between high subjective confidence and high objective performance, and vice versa. No correspondence between objective performance and subjective confidence was found in OCD ($R = 0.11$, $p = 0.67$) (online Supplementary Table S5.1 and online Supplementary Fig. S5.2).

ERP

Our data-driven windows for the HEP (150–320 ms) and the MP (–50 to 50 ms) resemble the canonical ones for both components (Deecke *et al.* 1984; Montoya *et al.* 1993; Johannes *et al.* 2001; Leopold & Schandry, 2001; Caldara *et al.* 2004; Pollatos & Schandry, 2004; Aravena *et al.* 2010).

HEP

First, to test the influence of cardiac attention on the HEP (Montoya *et al.* 1993; Pollatos *et al.* 2005), we




Fig. 1. Heartbeat detection task and associated heart evoked potential (HEP) modulations. (a) Behavioral results. Obsessive-compulsive disorder (OCD) obtained lower tapping accuracy (TA) scores, which indicates higher interoceptive accuracy (IAcc) than controls ($p < 0.01$) and panic patients ($p < 0.01$) (see online Supplementary Table S5.1 for details). The y axis indicates TA (in s). Values are means, with standard deviations represented by vertical bars. * Significant differences ($p < 0.05$). (b) HEP results. More negative HEP modulations for OCD in the exteroceptive (top) and interoceptive (bottom) conditions. In the exteroceptive condition, OCD patients showed a significantly more negative HEP than controls [latency = 303 ms, $t = 36.49$, degrees of freedom (df) = 12.46, $p < 0.01$] and panic patients (latency = 262 ms, $t = 29.77$, df = 91.62, $p < 0.01$). The same was true of the interoceptive condition, where OCD patients also showed a larger HEP than controls (latency = 190 ms, $t = 27.04$, df = 87.63, $p < 0.01$) and panic patients (latency = 182 ms, $t = 28.01$, df = 11.4, $p < 0.01$). Shaded bars around potentials indicate standard errors of the mean. * Significant difference between OCD and controls ($p < 0.05$). * Significant difference between OCD and panic ($p < 0.05$). No differences were found in either condition between controls and panic patients.

compared its modulation during the interoceptive condition against an exteroceptive one and a 7-min resting-state condition across groups. Controls showed more negative HEP amplitude in the interoceptive relative to both the exteroceptive and the resting-state conditions. The same modulation was observed in OCD, but no differences emerged in the panic sample (see online Supplementary material section 6.1 and Supplementary Fig. S6.1). While the latter is an interesting finding, it falls outside the main focus of our study; thus, we address it in online Supplementary material section 7.

A between-group comparison revealed significantly larger HEP amplitudes in OCD patients than in both other samples across conditions (Fig. 1b). As shown through a point-by-point comparison along the ERP via the permutation test, these differences emerged in the following expected windows (Leopold & Schandry, 2001; Fukushima et al. 2011; Shao et al. 2011): 200–300 ms for the exteroceptive condition,

and 160–230 ms for the interoceptive condition. No differences were found between controls and panic patients. Based on other studies that reported HEP modulations after the 500-ms mark, we replicated our analysis including epochs from -50 to 800 ms, and found the same pattern of increased amplitude in OCD patients in later windows (see online Supplementary Fig. S6.2).

MP

In the exteroceptive condition, the MP was more negative for correct than incorrect trials across groups (Fig. 2a). However, in line with previous research (Johannes et al. 2001; Ibanez et al. 2013), this modulation was more negative and wider (from -70 to 50 ms) in OCD patients. In panic patients, significant differences among categories emerged in a smaller window (from 40 to 70 ms). Between-group differences in these MP modulations are shown upon subtracting




Fig. 2. Motor potential (MP) modulations. (a) Intra-group comparisons: MP modulations differed between correct and incorrect responses in both conditions for obsessive-compulsive disorder (OCD) patients [exteroceptive condition: latency = 0.05 ms, $t = 55.45$, degrees of freedom (df) = 19.90, $p < 0.01$; interoceptive condition: latency = -0.04 ms, $t = 33.51$, df = 10.63, $p < 0.01$]. Differences in the exteroceptive condition were also observed in controls (latency = 0.03 ms, $t = 20.31$, df = 33.41, $p < 0.01$) and panic patients (latency = 0.08 ms, $t = 20.36$, df = 17.14, $p < 0.01$). The interoceptive condition yielded no significant results in panic patients and significant differences in only one point of the event-related potential for controls (latency = 0.8 ms; $t = 25.28$, df = 32.68, $p < 0.01$). (b) Inter-group comparisons. The subtraction between correct and incorrect responses showed a more negative amplitude of the MP in both conditions for OCD patients compared with controls and panic patients (exteroceptive condition: OCD > controls: latency = 0.03 ms, $t = 46.69$, df = 19.12, $p < 0.01$; OCD > panic: latency = 0.01 ms, 50.28, df = 18.21, $p < 0.01$; interoceptive condition: OCD > controls: latency = 0.05 ms, $t = 42.04$, df = 20.41, $p < 0.01$; OCD > panic: latency = 0.05 ms, $t = 46.41$, df = 13.69, $p < 0.01$). Shaded bars around potentials indicate standard errors of the mean. * Significant difference between OCD and controls ($p < 0.05$). * Significant difference between OCD and panic ($p < 0.05$). No differences were found between controls and panic patients.

incorrect from correct trials (Fig. 2b): more negative amplitudes were found in OCD patients relative to the other two groups, which showed similar MP amplitudes.

Additional differences were observed in the interoceptive condition (Fig. 2a). A larger negative amplitude for correct than incorrect trials was observed only in OCD patients. This pattern emerged in a window between -120 and -50 ms. The healthy control group showed a small difference (only one point of the ERP) in the same direction. No differences were found in the panic control sample. To gain further insights into these differential modulations, we compared the correct–incorrect difference wave between groups (Fig. 2b). The OCD sample exhibited a significantly larger modulation than the other two groups in the window from -80 to 20 ms. In sum, the difference in MP modulations between correct and incorrect responses was significantly higher in the OCD group than in the other two samples.

Interoception and dysfunctional beliefs in OCD

Using multiple regression models, we found that only the perfectionism subscale from the OBQ-TRIP was associated with the predictors mentioned in the Data analysis section ($R^2 = 0.48$, $p < 0.01$). More particularly, this association was significant relative to HEP modulations in the 300 – 400 ms window ($\beta = -0.81$, $p < 0.01$), but it was not significant relative to IAcc scores ($\beta = 0.21$, $p = 0.40$). No other significant associations were found (see online Supplementary Fig. S4 and Supplementary Table S4).

Discussion

By combining behavioral and electrophysiological measures, we evaluated the scope and distinctiveness of overactive monitoring in OCD. Patients with this condition showed increased IAcc and deficits in metacognitive abilities relative to controls and panic patients. Moreover, as compared with these two samples, they exhibited enhanced negativity of interoceptively relevant ERPs. This suggests the existence of condition-distinctive neural markers of altered inner monitoring in OCD. Below we discuss our key findings separately.

The scope of self-monitoring alterations in OCD

Previous behavioral studies failed to find increased monitoring in pediatric OCD samples (Hajcak et al. 2008; Hanna et al. 2012) and healthy but obsessive adults (Kaczurkin, 2013). However, the behavioral performance of OCD patients in our study was superior to that of controls, specifically during the

interoceptive condition of the HBD task. Together with evidence of fewer errors in flanker tasks (Riesel et al. 2011, 2014), our results add to the empirical corpus of enhanced monitoring accuracy in this population. Moreover, they highlight the sensitivity of cardiac monitoring tasks to address the issue.

Yet, as was the case in previous research (Endrass et al. 2008; Riesel et al. 2014), the most robust evidence of altered monitoring was obtained through electrophysiological measures. OCD patients consistently exhibited enhanced HEP and MP negativity during both external and internal monitoring. Although there are some discrepancies relative to the positive or negative deflection of the HEP, studies with parameters similar to our own have also reported negative deflections of this ERP (for a more detailed treatment of this, see online Supplementary material section 8). Thus, not only have we replicated this population's enhanced monitoring of environmental stimuli (Gehring et al. 2000; Riesel et al. 2011), but we have also shown that this pattern extends to inner monitoring skills.

In fact, interoceptive monitoring, as indexed by the MP, seems to be even more enhanced than its exteroceptive counterpart in OCD. Although MP modulations between correct and incorrect responses were observed in all groups, only OCD patients showed such an effect in the interoceptive condition. Moreover, their behavioral advantages were observed only when cardiac signals were triggered internally.

This pattern of inner overmonitoring may reflect biofunctional mechanisms associated with obsessive-compulsive cognition and symptoms. Indeed, inner monitoring (as indexed by HEP modulations) correlated with perfectionism ratings in OCD patients. Perfectionism is a core cognitive trait in OCD (Salkovskis, 1985; Coles et al. 2003; Belloch et al. 2010; Schrijvers et al. 2010) and it has been linked to the 'just right or not just right' (NJR) sensation (Coles et al. 2003; Starcevic et al. 2011). The latter has been described as a sensory or inner-drive feeling that disrupts homeostasis and prompts compulsive behaviors until things are 'right or perfect' (Coles et al. 2003) or until they reach a sense of completeness (Salkovskis et al. 2000). It has been suggested that this difficulty to stop can also explain the prolonged periods of mental rituals that patients with OCD perform to control their unwanted thoughts. Cognitive theories of obsessions state that patients with OCD struggle to decide whether they were successful at removing intrusive distressing thoughts (Salkovskis, 1985). This persistent doubt triggers either mental or behavioral rituals aimed at ensuring certainty that they do not hold any threatening thought (e.g. a patient who prays repeatedly to dispel blasphemous ideas). Thus, OCD

patients with high levels of perfectionism might present an attentional bias to bodily signals, which might sustain or even increase this NJR sensation. Such a conjecture aligns with evidence of an association between this cognitive domain and body checking in OCD (Vartanian & Germeroth, 2011). Hence, the alteration in the monitoring of bodily signals in OCD may represent an important mechanism affecting cognitive traits like perfectionism and associated symptoms such as checking behaviors, symmetry and ordering.

Although preliminary (due to the patients' sample size and the lack of a division according to OCD subtypes), our results open a novel avenue to assess the relationship between inner monitoring and cognitive/behavioral hallmarks of OCD. For example, patients who show fear reduction and become less hypervigilant about their obsessions after response prevention (EX/RP) treatment could also show a profile of reduced inner monitoring that could be tapped via HEP analysis. Future studies might include HEP modulations as a dependent measure in clinical trials to further our understanding of brain changes after EX/RP treatment.

Metacognition of inner over-monitoring in OCD

The OCD patients were less confident about their actual performance than the other samples. This was true despite their objectively superior performance. Such a dissociation between objective performance and subjective confidence was further supported by IAW results. Indeed, the OCD patients were the only ones showing no correspondence between objective performance and subjective estimation of their outcome.

This finding aligns with previous reports showing metacognitive deficits in OCD (Cucchi *et al.* 2012; Coles *et al.* 2015), which are acknowledged as core symptoms in psychopathological models of the disorder (Hermans *et al.* 2008). Compatibly, our results suggest that classical metacognitive alterations relative to external monitoring (Endrass *et al.* 2008, 2010) also extend to the monitoring of inner bodily signals. Unlike previous metacognition results showing no differences between OCD and other clinical samples (Janeck *et al.* 2003; Cucchi *et al.* 2012; Brevers *et al.* 2013; Odlaug *et al.* 2014), our findings suggest that OCD could be characterized by alterations in a specific metacognitive subskill: the ability to estimate one's own internal attentional abilities.

The 'specificity' of self-monitoring alterations in OCD

While the external overmonitoring pattern found in OCD has also been observed in patients with general

anxiety (Stern *et al.* 2010; Xiao *et al.* 2011) and post-traumatic stress disorders (Rabinak *et al.* 2013), research on the specificity of self-monitoring alterations in OCD has so far yielded inconclusive results (Gehring *et al.* 2000; Endrass *et al.* 2008; Weinberg *et al.* 2010; Riesel *et al.* 2011, 2014; Xiao *et al.* 2011; Carrasco *et al.* 2013; Moser *et al.* 2013; Lin *et al.* 2015). The same is true of electrophysiological studies on processing external stimulus monitoring: whereas some studies have shown higher ERN and CRN amplitude in OCD patients compared with general anxiety disorder patients (Carrasco *et al.* 2013), others yielded the opposite pattern (Xiao *et al.* 2011).

However, no study has assessed inner monitoring in OCD via interoceptive tasks. Previous findings in other anxiety disorders are inconclusive, with patients performing either better than (Ehlers & Breuer, 1992), worse than (Willem Van der Does *et al.* 2000) or similar to (Yoris *et al.* 2015) controls. Despite this controversy, these reports highlight that patients with anxiety disorders might not present an overmonitoring of bodily signals as we found in our OCD sample. Also, no study seems to have assessed HEP modulations in OCD solely, or in comparison with other pathological conditions. However, there are a few studies that have found reduced HEP modulation in other psychiatric conditions, such as depression (Terhaar *et al.* 2012), borderline personality disorder (Muller *et al.* 2015) and depersonalization disorder (Schulz *et al.* 2015). These findings suggest that inner overmonitoring alterations, as indexed by IAcc scores and HEP modulations, might represent a distinctive hallmark of OCD patients.

In sum, focusing on interoceptive monitoring, we found significant differences between OCD and panic disorder patients across behavioral and electrophysiological dimensions. Specifically, OCD patients exhibited: (i) more accurate performance in the interoceptive than the exteroceptive condition; (ii) more negative HEP and MP amplitude in both conditions; (iii) earlier modulations of these components in the interoceptive condition; and (iv) lower ISen and IAW scores. Thus, cross-dimensional aspects of internal monitoring seem distinctively altered in OCD.

Given that OCD shares co-morbid symptoms with other psychiatric disorders (e.g. 75.6% with any other anxiety disorder, 40.7% with major mood disorder, and 20% with panic; Ruscio *et al.* 2010), the apparent specificity of interoceptive alterations across the anxiety spectrum may offer a new, valuable contribution for accurate diagnosis and longitudinal tracking of this condition. Thus, in line with evidence of differential neural markers of symptom groups in OCD (Mataix-Cols *et al.* 2004), our results suggest that psychiatric assessment may be fruitfully complemented

with behavioral and neurocognitive assessment of target domains. Moreover, by evincing a differential link between cognitive and bodily functions, they contribute to forging post-dualistic approaches to the study of psychiatric conditions (Zinbarg & Barlow, 1996; Barlow & Campbell, 2000).

Limitations and avenues for further research

Our sample size was modest. However, previous studies with similar numbers of participants also reported robust results (Gehring et al. 2000; Johannes et al. 2001; Endrass et al. 2008; Stern et al. 2010). Besides, we were unable to consider different subtypes of OCD. Yet, all our patients featured similar symptoms and each of them had OCI-R scores two standard deviations above those of the panic group, which speaks to the sample's broad homogeneity. That being said, since this disorder features high symptomatic variability (Foa et al. 2002), future research should favor a dimensional approach comparing self-monitoring patterns across subtypes of OCD. In addition, although the inclusion of panic patients represents an initial effort to evaluate the specificity of inner overmonitoring alterations in OCD, future studies should include samples featuring other anxiety disorders to more rigorously test this possibility.

Finally, different interoceptive accuracy tasks have been used in the literature (e.g. the discrimination task and the mental tracking task), each with its own advantages and disadvantages (Domschke et al. 2010; Yoris et al. 2015). We selected an interoceptive task that requires motor tapping because external overmonitoring in OCD is generally tested with motor tasks (Endrass & Ullsperger, 2014; Riesel et al. 2014, 2015). Importantly, even if one assumes that a motor confound exists in the results of the OCD group, behavioral differences between groups should have emerged across conditions rather than only in the interoceptive task. Also, in the latter, OCD patients exhibited greater IAcc than the other groups, which was coherent with HEP differences. Finally, previous studies have also shown that OCD performance in motor tasks varies widely, from good to bad (Gehring et al. 2000). Still, a replication of our study with non-motor interoceptive tasks could offer useful insights on the issue.

Conclusion

Our work offers unprecedented cross-dimensional evidence of altered interoceptive over-monitoring in OCD. Convergent behavioral and electrophysiological data showed that overactive monitoring in OCD extends to the sensing of internal bodily signals. Notably, this pattern discriminated OCD from controls

and panic patients, suggesting a condition-distinctive alteration. These results highlight the potential of exploring interoceptive processes in the OCD spectrum to: (i) better characterize the population's cognitive profile; (ii) evaluate their sensitivity and specificity as a potential endophenotype (Riesel et al. 2015); (iii) assess their relation with characteristic obsessive-compulsive cognitive profiles and symptoms; and (iv) evaluate their potential relevance to track disease progression, therapeutic impact and other forms of clinical response. Finally, our findings pave the way for new insights into the links between overmonitoring, cognition and body sensing. This is the first study to show that one of the most replicated hallmarks of OCD seems to extend to the body's inner world, opening new fields of research and assessment.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717000368>

Acknowledgements

This work was partially supported by grants from CONICET, CONICYT/FONDECYT Regular (1130920 and 1140114), FONCYT-PICT 2012-0412, FONCYT-PICT 2012-1309, FONDAP 15150012, and the INECO Foundation.

Declaration of Interest

None.

References

- Aouizerate B, Guehl D, Cuny E, Rougier A, Bioulac B, Tignol J, Burbaud P (2004). Pathophysiology of obsessive-compulsive disorder: a necessary link between phenomenology, neuropsychology, imagery and physiology. *Progress in Neurobiology* **72**, 195–221.
- American Psychiatric Association (1994). *DSM-IV: Diagnostic and Statistical Manual*. American Psychiatric Association: Washington, DC.
- Aravena P, Hurtado E, Riveros R, Cardona JF, Manes F, Ibanez A (2010). Applauding with closed hands: neural signature of action-sentence compatibility effects. *PLoS ONE* **5**, e11751.
- Bannon S, Gonsalvez CJ, Croft RJ, Boyce PM (2002). Response inhibition deficits in obsessive-compulsive disorder. *Psychiatry Research* **110**, 165–174.
- Barlow DH, Campbell LA (2000). Mixed anxiety-depression and its implications for models of mood and anxiety disorders. *Comprehensive Psychiatry* **41**, 55–60.
- Belloch A, Morillo C, Luciano JV, Garcia-Soriano G, Cabedo E, Carrio C (2010). Dysfunctional belief domains related to obsessive-compulsive disorder: a further examination of

- their dimensionality and specificity. *Spanish Journal of Psychology* **13**, 376–388.
- Brener J, Kluvitse C** (1988). Heartbeat detection: judgments of the simultaneity of external stimuli and heartbeats. *Psychophysiology* **25**, 554–561.
- Brevers D, Cleeremans A, Bechara A, Greisen M, Kornreich C, Verbanck P, Noel X** (2013). Impaired self-awareness in pathological gamblers. *Journal of Gambling Studies* **29**, 119–129.
- Brunia CH, Van den Bosch WE** (1984). Movement-related slow potentials. I. A contrast between finger and foot movements in right-handed subjects. *Electroencephalography and Clinical Neurophysiology* **57**, 515–527.
- Caldara R, Deiber MP, Andrey C, Michel CM, Thut G, Hauert CA** (2004). Actual and mental motor preparation and execution: a spatiotemporal ERP study. *Experimental Brain Research* **159**, 389–399.
- Cameron OG** (2001). Interoception: the inside story – a model for psychosomatic processes. *Psychosomatic Medicine* **63**, 697–710.
- Canales-Johnson A, Silva C, Huepe D, Rivera-Rei A, Noreika V, Garcia MD, Silva W, Ciraolo C, Vaucheret E, Sedeno L, Couto B, Kargieman L, Baglivo F, Sigman M, Chennu S, Ibanez A, Rodriguez E, Bekinschtein TA** (2015). Auditory feedback differentially modulates behavioral and neural markers of objective and subjective performance when tapping to your heartbeat. *Cerebral Cortex* **25**, 4490–4503.
- Carrasco M, Harbin SM, Nienhuis JK, Fitzgerald KD, Gehring WJ, Hanna GL** (2013). Increased error-related brain activity in youth with obsessive–compulsive disorder and unaffected siblings. *Depression and Anxiety* **30**, 39–46.
- Coles ME, Frost RO, Heimberg RG, Rheume J** (2003). “Not just right experiences”: perfectionism, obsessive–compulsive features and general psychopathology. *Behaviour Research and Therapy* **41**, 681–700.
- Coles ME, Schofield CA, Nota JA** (2015). Initial data on recollections of pathways to inflated responsibility beliefs in patients with obsessive–compulsive disorder. *Behavioural and Cognitive Psychotherapy* **43**, 385–395.
- Couto B, Adolff F, Velasquez M, Mesow M, Feinstein J, Canales-Johnson A, Mikulan E, Martinez-Pernia D, Bekinschtein T, Sigman M, Manes F, Ibanez A** (2015). Heart evoked potential triggers brain responses to natural affective scenes: a preliminary study. *Autonomic Neuroscience: Basic and Clinical* **193**, 132–137.
- Couto B, Salles A, Sedeno L, Peradejordi M, Barttfeld P, Canales-Johnson A, Dos Santos YV, Huepe D, Bekinschtein T, Sigman M, Favaloro R, Manes F, Ibanez A** (2014). The man who feels two hearts: the different pathways of interoception. *Social Cognitive and Affective Neuroscience* **9**, 1253–1260.
- Craig AD** (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews. Neuroscience* **3**, 655–666.
- Craig AD** (2003). Interoception: the sense of the physiological condition of the body. *Current Opinion in Neurobiology* **13**, 500–505.
- Craig AD** (2004). Human feelings: why are some more aware than others? *Trends in Cognitive Sciences* **8**, 239–241.
- Critchley HD, Harrison NA** (2013). Visceral influences on brain and behavior. *Neuron* **77**, 624–638.
- Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ** (2004). Neural systems supporting interoceptive awareness. *Nature Neuroscience* **7**, 189–195.
- Cucchi M, Bottelli V, Cavadini D, Ricci L, Conca V, Ronchi P, Smeraldi E** (2012). An explorative study on metacognition in obsessive–compulsive disorder and panic disorder. *Comprehensive Psychiatry* **53**, 546–553.
- Deecke L, Bashore T, Brunia CH, Grunewald-Zuberbier E, Grunewald G, Kristeva R** (1984). Movement-associated potentials and motor control. Report of the EPIC VI Motor Panel. *Annals of the New York Academy of Sciences* **425**, 398–428.
- Deecke L, Eisinger H, Kornhuber HH** (1980). Comparison of Bereitschaftspotential, pre-motion positivity and motor potential preceding voluntary flexion and extension movements in man. *Progress in Brain Research* **54**, 171–176.
- Domschke K, Stevens S, Pfeleiderer B, Gerlach AL** (2010). Interoceptive sensitivity in anxiety and anxiety disorders: an overview and integration of neurobiological findings. *Clinical Psychology Review* **30**, 1–11.
- Dum RP, Levinthal DJ, Strick PL** (2009). The spinothalamic system targets motor and sensory areas in the cerebral cortex of monkeys. *Journal of Neuroscience: the Official Journal of the Society for Neuroscience* **29**, 14223–14235.
- Dunn BD, Dalglish T, Ogilvie AD, Lawrence AD** (2007). Heartbeat perception in depression. *Behaviour Research and Therapy* **45**, 1921–1930.
- Ehlers A, Breuer P** (1992). Increased cardiac awareness in panic disorder. *Journal of Abnormal Psychology* **101**, 371–382.
- Endrass T, Klawohn J, Schuster F, Kathmann N** (2008). Overactive performance monitoring in obsessive–compulsive disorder: ERP evidence from correct and erroneous reactions. *Neuropsychologia* **46**, 1877–1887.
- Endrass T, Schuermann B, Kaufmann C, Spielberg R, Kniesche R, Kathmann N** (2010). Performance monitoring and error significance in patients with obsessive–compulsive disorder. *Biological Psychology* **84**, 257–263.
- Endrass T, Ullsperger M** (2014). Specificity of performance monitoring changes in obsessive–compulsive disorder. *Neuroscience and Biobehavioral Reviews* **46**, 124–138.
- First MB, Spitzer RL, Gibbon M, Williams JB** (2002). *Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Patient Edition (SCID-I/P)*. Biometrics Research, New York State Psychiatric Institute: New York.
- Foa EB, Huppert JD, Leiberg S, Langner R, Kichic R, Hajcak G, Salkovskis PM** (2002). The Obsessive–Compulsive Inventory: development and validation of a short version. *Psychological Assessment* **14**, 485–496.
- Fukushima H, Terasawa Y, Umeda S** (2011). Association between interoception and empathy: evidence from heartbeat-evoked brain potential. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology* **79**, 259–265.
- García-Cordero I, Sedeño L, de la Fuente L, Slachevsky A, Forno G, Klein F, Lillo P, Ferrari J, Rodriguez C, Bustin J, Torralva T, Baez S, Yoris A, Esteves S, Melloni M, Salamone P, Huepe D, Manes F, García AM, Ibañez A**

- (2016). Feeling, learning from, and being aware of inner states: interoceptive dimensions in neurodegeneration and stroke. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **371**, 20160006.
- Garfinkel SN, Seth AK, Barrett AB, Suzuki K, Critchley HD** (2015). Knowing your own heart: distinguishing interoceptive accuracy from interoceptive awareness. *Biological Psychology* **104**, 65–74.
- Gehring WJ, Himle J, Nisenson LG** (2000). Action-monitoring dysfunction in obsessive–compulsive disorder. *Psychological Science* **11**, 1–6.
- Gray MA, Taggart P, Sutton PM, Groves D, Holdright DR, Bradbury D, Brull D, Critchley HD** (2007). A cortical potential reflecting cardiac function. *Proceedings of the National Academy of Sciences USA* **104**, 6818–6823.
- Hajcak G, Franklin ME, Foa EB, Simons RF** (2008). Increased error-related brain activity in pediatric obsessive–compulsive disorder before and after treatment. *American Journal of Psychiatry* **165**, 116–123.
- Hanna GL, Carrasco M, Harbin SM, Nienhuis JK, LaRosa CE, Chen P, Fitzgerald KD, Gehring WJ** (2012). Error-related negativity and tic history in pediatric obsessive–compulsive disorder. *Journal of the American Academy of Child and Adolescent Psychiatry* **51**, 902–910.
- Hermans D, Engelen U, Grouwels L, Joos E, Lemmens J, Pieters G** (2008). Cognitive confidence in obsessive–compulsive disorder: distrusting perception, attention and memory. *Behaviour Research and Therapy* **46**, 98–113.
- Ibanez A, Cardona JF, Dos Santos YV, Blenkmann A, Aravena P, Roca M, Hurtado E, Nerguizian M, Amoroso L, Gomez-Arevalo G, Chade A, Dubrovsky A, Gershanik O, Kochen S, Glenberg A, Manes F, Bekinschtein T** (2013). Motor–language coupling: direct evidence from early Parkinson’s disease and intracranial cortical recordings. *Cortex* **49**, 968–984.
- Janeck AS, Calamari JE, Riemann BC, Heffelfinger SK** (2003). Too much thinking about thinking?: metacognitive differences in obsessive–compulsive disorder. *Journal of Anxiety Disorders* **17**, 181–195.
- Janig W** (1996). Neurobiology of visceral afferent neurons: neuroanatomy, functions, organ regulations and sensations. *Biological Psychology* **42**, 29–51.
- Johannes S, Wieringa BM, Mantey M, Nager W, Rada D, Muller-Vahl KR, Emrich HM, Dengler R, Munte TF, Dietrich D** (2001). Altered inhibition of motor responses in Tourette syndrome and obsessive–compulsive disorder. *Acta Neurologica Scandinavica* **104**, 36–43.
- Kaczurkin AN** (2013). The effect of manipulating task difficulty on error-related negativity in individuals with obsessive–compulsive symptoms. *Biological Psychology* **93**, 122–131.
- Khalsa SS, Rudrauf D, Feinstein JS, Tranel D** (2009). The pathways of interoceptive awareness. *Nature Neuroscience* **12**, 1494–1496.
- Kleckner IR, Wormwood JB, Simmons WK, Barrett LF, Quigley KS** (2015). Methodological recommendations for a heartbeat detection-based measure of interoceptive sensitivity. *Psychophysiology* **52**, 1432–1440.
- Kruczyk M, Umer HM, Enroth S, Komorowski J** (2013). Peak finder metaserver – a novel application for finding peaks in ChIP-seq data. *BMC Bioinformatics* **14**, 280.
- Leopold C, Schandry R** (2001). The heartbeat-evoked brain potential in patients suffering from diabetic neuropathy and in healthy control persons. *Clinical Neurophysiology* **112**, 674–682.
- Lin Y, Moran TP, Schroder HS, Moser JS** (2015). The role of hand of error and stimulus orientation in the relationship between worry and error-related brain activity: implications for theory and practice. *Psychophysiology* **52**, 1281–1292.
- Manly BFJ** (2007). *Randomization, Bootstrap, and Monte Carlo Methods in Biology*. Chapman & Hall/CRC: Boca Raton, FL.
- Mataix-Cols D, Wooderson S, Lawrence N, Brammer MJ, Speckens A, Phillips ML** (2004). Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessive–compulsive disorder. *Archives of General Psychiatry* **61**, 564–576.
- Montoya P, Schandry R, Muller A** (1993). Heartbeat evoked potentials (HEP): topography and influence of cardiac awareness and focus of attention. *Electroencephalography and Clinical Neurophysiology* **88**, 163–172.
- Moser JS, Moran TP, Schroder HS, Donnellan MB, Yeung N** (2013). On the relationship between anxiety and error monitoring: a meta-analysis and conceptual framework. *Frontiers in Human Neuroscience* **7**, 466.
- Moulding R, Anglim J, Nedeljkovic M, Doron G, Kyrios M, Ayalon A** (2011). The Obsessive Beliefs Questionnaire (OBQ): examination in nonclinical samples and development of a short version. *Assessment* **18**, 357–374.
- Muller LE, Schulz A, Andermann M, Gabel A, Gescher DM, Spohn A, Herpertz SC, Bertsch K** (2015). Cortical representation of afferent bodily signals in borderline personality disorder: neural correlates and relationship to emotional dysregulation. *JAMA Psychiatry* **72**, 1077–1086.
- Nagai M, Kishi K, Kato S** (2007). Insular cortex and neuropsychiatric disorders: a review of recent literature. *European Psychiatry: the Journal of the Association of European Psychiatrists* **22**, 387–394.
- Nichols TE, Holmes AP** (2002). Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Human Brain Mapping* **15**, 1–25.
- Odlaug BL, Chamberlain SR, Derbyshire KL, Leppink EW, Grant JE** (2014). Impaired response inhibition and excess cortical thickness as candidate endophenotypes for trichotillomania. *Journal of Psychiatric Research* **59**, 167–173.
- Pollatos O, Herbert BM, Mai S, Kammer T** (2016). Changes in interoceptive processes following brain stimulation. *Philosophical Transaction of the Royal Society of London* **371**, 20160016.
- Pollatos O, Kirsch W, Schandry R** (2005). Brain structures involved in interoceptive awareness and cardioafferent signal processing: a dipole source localization study. *Human Brain Mapping* **26**, 54–64.
- Pollatos O, Matthias E, Schandry R** (2007). Heartbeat perception and P300 amplitude in a visual oddball paradigm. *Clinical Neurophysiology* **118**, 2248–2253.

- Pollatos O, Schandry R (2004). Accuracy of heartbeat perception is reflected in the amplitude of the heartbeat-evoked brain potential. *Psychophysiology* **41**, 476–482.
- Rabinak CA, Holman A, Angstadt M, Kennedy AE, Hajcak G, Phan KL (2013). Neural response to errors in combat-exposed returning veterans with and without post-traumatic stress disorder: a preliminary event-related potential study. *Psychiatry Research* **213**, 71–78.
- Riesel A, Endrass T, Auerbach LA, Kathmann N (2015). Overactive performance monitoring as an endophenotype for obsessive-compulsive disorder: evidence from a treatment study. *American Journal of Psychiatry* **172**, 665–673.
- Riesel A, Endrass T, Kaufmann C, Kathmann N (2011). Overactive error-related brain activity as a candidate endophenotype for obsessive-compulsive disorder: evidence from unaffected first-degree relatives. *American Journal of Psychiatry* **168**, 317–324.
- Riesel A, Kathmann N, Endrass T (2014). Overactive performance monitoring in obsessive-compulsive disorder is independent of symptom expression. *European Archives of Psychiatry and Clinical Neuroscience* **264**, 707–717.
- Ronchi R, Bello-Ruiz J, Lukowska M, Herbelin B, Cabrilo I, Schaller K, Blanke O (2015). Right insular damage decreases heartbeat awareness and alters cardio-visual effects on bodily self-consciousness. *Neuropsychologia* **70**, 11–20.
- Ruscio AM, Stein DJ, Chiu WT, Kessler RC (2010). The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Molecular Psychiatry* **15**, 53–63.
- Salkovskis PM (1985). Obsessional-compulsive problems: a cognitive-behavioural analysis. *Behaviour Research and Therapy* **23**, 571–583.
- Salkovskis PM, Wroe AL, Gledhill A, Morrison N, Forrester E, Richards C, Reynolds M, Thorpe S (2000). Responsibility attitudes and interpretations are characteristic of obsessive compulsive disorder. *Behaviour Research and Therapy* **38**, 347–372.
- Schandry R, Bestler M, Montoya P (1993). On the relation between cardiodynamics and heartbeat perception. *Psychophysiology* **30**, 467–474.
- Schandry R, Sparrer B, Weitkunat R (1986). From the heart to the brain: a study of heartbeat contingent scalp potentials. *International Journal of Neuroscience* **30**, 261–275.
- Schandry R, Weitkunat R (1990). Enhancement of heartbeat-related brain potentials through cardiac awareness training. *International Journal of Neuroscience* **53**, 243–253.
- Schrijvers DL, De Bruijn ER, Destoop M, Hulstijn W, Sabbe BG (2010). The impact of perfectionism and anxiety traits on action monitoring in major depressive disorder. *Journal of Neural Transmission (Vienna)* **117**, 869–880.
- Schulz A, Koster S, Beutel ME, Schachinger H, Vogele C, Rost S, Rauh M, Michal M (2015). Altered patterns of heartbeat-evoked potentials in depersonalization/derealization disorder: neurophysiological evidence for impaired cortical representation of bodily signals. *Psychosomatic Medicine* **77**, 506–516.
- Schulz A, Matthey JH, Vogele C, Schaan V, Schachinger H, Adler J, Beutel ME, Michal M (2016). Cardiac modulation of startle is altered in depersonalization-/derealization disorder: evidence for impaired brainstem representation of baro-afferent neural traffic. *Psychiatry Research* **240**, 4–10.
- Sedeno L, Couto B, Melloni M, Canales-Johnson A, Yoris A, Baez S, Esteves S, Velasquez M, Barttfeld P, Sigman M, Kichic R, Chialvo D, Manes F, Bekinschtein TA, Ibanez A (2014). How do you feel when you can't feel your body? Interoception, functional connectivity and emotional processing in depersonalization-derealization disorder. *PLOS ONE* **9**, e98769.
- Shao S, Shen K, Wilder-Smith EP, Li X (2011). Effect of pain perception on the heartbeat evoked potential. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology* **122**, 1838–1845.
- Simons RF (2010). The way of our errors: theme and variations. *Psychophysiology* **47**, 1–14.
- Song A, Jung WH, Jang JH, Kim E, Shim G, Park HY, Choi CH, Kwon JS (2011). Disproportionate alterations in the anterior and posterior insular cortices in obsessive-compulsive disorder. *PLoS ONE* **6**, e22361.
- Spielberger CD, Gorsuch RL, Lushene RE (1970). *Manual for the State-Trait Anxiety Inventory*. Consulting Psychological Press: Palo Alto, CA.
- Starcevic V, Berle D, Brakoulias V, Sammut P, Moses K, Milicevic D, Hannan A (2011). Functions of compulsions in obsessive-compulsive disorder. *Australian and New Zealand Journal of Psychiatry* **45**, 449–457.
- Stern ER, Liu Y, Gehring WJ, Lister JJ, Yin G, Zhang J, Fitzgerald KD, Himle JA, Abelson JL, Taylor SF (2010). Chronic medication does not affect hyperactive error responses in obsessive-compulsive disorder. *Psychophysiology* **47**, 913–920.
- Terasawa Y, Kurosaki Y, Ibata Y, Moriguchi Y, Umeda S (2015). Attenuated sensitivity to the emotions of others by insular lesion. *Frontiers in Psychology* **6**, 1314.
- Terhaar J, Viola FC, Bar KJ, Debener S (2012). Heartbeat evoked potentials mirror altered body perception in depressed patients. *Clinical Neurophysiology* **123**, 1950–1957.
- Vartanian LR, Germeroth LJ (2011). Accuracy in estimating the body weight of self and others: impact of dietary restraint and BMI. *Body Image* **8**, 415–418.
- Vaughan Jr. HG, Costa LD, Ritter W (1968). Topography of the human motor potential. *Electroencephalography and Clinical Neurophysiology* **25**, 1–10.
- Weinberg A, Olvet DM, Hajcak G (2010). Increased error-related brain activity in generalized anxiety disorder. *Biological Psychology* **85**, 472–480.
- Weiss S, Sack M, Henningsen P, Pollatos O (2014). On the interaction of self-regulation, interoception and pain perception. *Psychopathology* **47**, 377–382.
- Wiens S, Palmer SN (2001). Quadratic trend analysis and heartbeat detection. *Biological Psychology* **58**, 159–175.
- Willem Van der Does AJ, Antony MM, Ehlers A, Barsky AJ (2000). Heartbeat perception in panic disorder: a reanalysis. *Behaviour Research and Therapy* **38**, 47–62.
- Xiao Z, Wang J, Zhang M, Li H, Tang Y, Wang Y, Fan Q, Fromson JA (2011). Error-related negativity abnormalities

in generalized anxiety disorder and obsessive–compulsive disorder. *Progress in Neuro-psychopharmacology and Biological Psychiatry* **35**, 265–272.

Yoris A, Esteves S, Couto B, Melloni M, Kichic R, Cetkovich M, Favaloro R, Moser J, Manes F, Ibanez A, Seden L (2015). The roles of interoceptive sensitivity and metacognitive interoception in panic. *Behavioral and Brain Functions: BBF* **11**, 14.

Zaki J, Davis JI, Ochsner KN (2012). Overlapping activity in anterior insula during interoception and emotional experience. *NeuroImage* **62**, 493–499.

Zigmond AS, Snaith RP (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica* **67**, 361–370.

Zinbarg RE, Barlow DH (1996). Structure of anxiety and the anxiety disorders: a hierarchical model. *Journal of Abnormal Psychology* **105**, 181–193.